

---

# Recent trends in summer atmospheric circulation in the North Atlantic/European region: is there a role for anthropogenic aerosols?

Rowan Sutton<sup>\*1</sup> and Buwen Dong<sup>†2</sup>

<sup>1</sup>National Centre for Atmospheric Science (Reading) – Department of Meteorology, University of Reading, PO Box 243, Earley Gate, Reading RG6 6BB., United Kingdom

<sup>2</sup>National Centre for Atmospheric Science – NCAS, Department of Meteorology, University of Reading, United Kingdom

## Abstract

The variability of the westerly jet stream and storm track is crucial for summer weather and climate in the North Atlantic/European region. Observations show notable trends in the summer jet from 1970s to 2010s, characterized by an equatorward migration over the North Atlantic and a poleward migration and weakening of the Mediterranean jet over Europe. These changes were associated with more cyclonic storms traveling across the UK, and fewer over the Mediterranean, leading to wet summers in northern Europe and dry summers in southern Europe. In this study we investigate the potential drivers and processes that may have been responsible for the observed changes in summer atmospheric circulation, with a particular focus on the role of anthropogenic aerosols (AA). We conduct attribution experiments with an atmospheric general circulation model forced with observed changes in sea surface temperatures/sea ice extent (SST/SIE), greenhouse gas concentrations and AA precursor emissions. Results strongly suggest that fast responses to AA changes were likely the primary driver of the observed poleward migration and weakening of the Mediterranean jet, with changes in SST/SIE playing a secondary role. The simulated response shows good agreement with the observed changes in both magnitude and vertical structure, which suggests that common mechanisms - involving aerosol-radiation and aerosol-cloud interactions - are responsible. By contrast, changes in the North Atlantic jet are influenced in the model experiments by changes in both Atlantic SST/SIE and fast responses to AA. In this case, however, there are significant differences between the model response and the observed changes, which may be explained by biases in the model climatology.

**Keywords:** atmospheric circulation, anthropogenic aerosols, jet stream, summer

---

<sup>\*</sup>Speaker

<sup>†</sup>Corresponding author: [buwen.dong@ncas.ac.uk](mailto:buwen.dong@ncas.ac.uk)